

POST-PROCEDURE CARE

After the LLETZ procedure you should not swim, soak in the bath or spa, use tampons or have vaginal intercourse for four to six weeks until the cervix has healed to avoid infection.

Strenuous exercise should also be avoided for seven to ten days as this increases the risk of bleeding.

You should seek medical advice if you experience any of the following:

- Persistent vaginal bleeding
- Passing large clots or tissue
- Offensive smelling vaginal discharge
- Temperature over 38°C, or chills
- Persistent nausea or vomiting
- Increasing abdominal not relieved by analgesia
- Pain, burning or frequency when passing urine
- Difficulty in emptying your bladder

If you have a concern post your procedure:

During Business Hours:

Contact WHC - 5594 9496

After Hours or Emergency:

GCUH or your nearest emergency department



**DR
DONALD
ANGSTETRA**

OBSTETRICIAN AND GYNAECOLOGIST
GYNAECOLOGY ENDOSCOPIC SURGEON

Women's Health Centre Gold Coast
Gold Coast Private Hospital
Suite 18, level 1, 14 Hill Street
Southport QLD 4215

reception@thewomenshealthcentre.com

P: 5594 9496 | F: 5594 9102

COLPOSCOPY & LLETZ

PATIENT INFORMATION